

---

# Web application for Intensity of Erosion and Outflow

**Name of the River Basin: Radulicka rijeka**

**Country: Montenegro**

**Year: 2018**

**GPS coordinates, latitude and longitude with Google Maps:  
42.965364,19.822026**

## INPUT DATA

### Geometric characteristics of the river basins

**F = 40.23096 km<sup>2</sup> (Surface area of the drainage basin)**

**O = 29.8793 km (Length of the watershed)**

**Fv = 27.78336 km<sup>2</sup> (Surface area of greater portion of the drainage basin)**

**Fm = 12.4476 km<sup>2</sup> (Surface area of smaller portion of the drainage basin)**

**Lv = 8.46906 km (Natural length of main water course)**

**Lb = 12.16551 km (Length of the drainage basin measured by a series of parallel lines)**

### Topographic characteristics of the river basins

**Contour line length - Liz [km]: ["20.14833 ", "33.17354 ", "30.38447 ", "14.58891 ", "9.80364 ", "7.63453 ", "2.80515 "]**

**The area between the two neighboring contour lines - f [km<sup>2</sup>]: ["4.88120 ", "9.81131 ", "9.50161 ", "7.45868 ", "3.61518 ", "2.96913 ", "1.60158 ", "0.39228 "]**

**h0 = 700 m (Altitude of the initial contour)**

**Δh = 100 m (Equidistance)**

**Hmin = 619 (Lowest altitude in the drainage basin)**

**Hmax = 1356 (Highest altitude in the drainage basin)**

---

## Hydrological characteristics of the river basins

$\Sigma L = 12.37249$  km (The total length of the main watercourse with tributaries of 1<sup>st</sup> and 2<sup>nd</sup> class)

$L_m = 7.0783$  km (The shortest distance between the fountain (head and mouth))

### Water permeability

$f_p = 0.1025$  (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0.1193$  (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.7782$  (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

### Land use

$f_s = 0.498684400$  (Part of the surface area of the drainage basin under the forest)

$f_t = 0.352763084$  (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.148552517$  (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

### Meteorological data

$h_b = 157.6$  mm (Level of torrent rain)

$U_p$  (years) = 100

$t_o = 8.9$  °C (Average annual air temperature)

$H_{god} = 983.7$  mm (Average annual quantity of precipitation)

### Erosion coefficients

$Y = 1.14102$  (Types of soil structures and allied types)

5.1 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

92.46 % (Serpentines, red sand stones, flishe deposits)

---

---

**0 % (Podzols and parapodzols, decomposed schist)**

**2.44 % (Solid and Schist limestone, Terra Rosa and Humic soil)**

**0 % (Brown forest soils and Mountain soils)**

**0 % (Epieugleysol and Marshlands)**

**0 % (Good structured Chernozems and alluvial well-structured deposits)**

**0 % (Bare, compact igneous)**

**Xa = 0.508745 (Planning of the drainage basin, rate of drainage basin regulation)**

**0 % (Bare lands)**

**14.86 % (Plough-lands)**

**1.89 % (Orchards and vineyards)**

**19.36 % (Mountain pastures)**

**14.03 % (Meadows)**

**29.92 % (Degraded forests)**

**19.95 % (Well-constituted forests)**

**$\phi = 0.31364$  (Numerical coefficient of visible and clearly pointed processes of soil erosion)**

**4.98 % (Depth erosion)**

**4.42 % (80% of the river basin under rill and gully erosion)**

**3.87 % (50% of the river basin under rill and gully erosion)**

**3.32 % (100% of the river basin under surface erosion)**

**14.03 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)**

**1.66 % (50% of the river basin under surface erosion)**

**1.11 % (20% of the river basin under surface erosion)**

**0 % (There are smaller slides in the watercourse beds)**

**14.86 % (The river basin mostly under plough-land)**

**51.76 % (The river basin under forests and perennial vegetation)**

## **INPUT DATA**

**A = 0.68797050676226 (Coefficient of the river basin form)**

**m = 0.376660500986 (Coefficient of the watershed development)**

**B = 3.3069686351004 km (Average river basin width)**

**a = 0.76238598333224 ((A)symmetry of the river basin)**

---

---

**G = 0.30753653405238 (Density of the river network of the basin)**  
**K = 1.1964822061795 (Coefficient of the river basin tortuousness)**  
**H<sub>sr</sub> = 881.73659639243 m (Average river basin altitude)**  
**D = 262.73659639243 m (Average elevation difference of the river basin)**  
**I<sub>sr</sub> = 29.46451439389 % (Average river basin decline)**  
**H<sub>leb</sub> = 737 m (The height of the local erosion base of the river basin)**  
**E<sub>r</sub> = 93.14885399898 (Coefficient of the erosion energy of the river basins relief)**  
**S<sub>1</sub> = 0.90271 (Coefficient of the regions permeability)**  
**S<sub>2</sub> = 0.7299736242 (Coefficient of the vegetation cover)**  
**W = 1.6799766814581 m (Analytical presentation of the water retention in inflow)**  
**2gDF<sup>1/2</sup> = 455.39681016669 m km s<sup>-1</sup> (Energetic potential of water flow during torrent rains)**  
**Q<sub>max</sub> = 346.83145682303 m<sup>3</sup> s<sup>-1</sup> (Maximal outflow from the river basin)**  
**T = 0.99498743710662 (Temperature coefficient of the region)**  
**Z = 0.49716044480634 (Coefficient of the river basin erosion)**  
**W<sub>god</sub> = 43364.604433765 m<sup>3</sup> god<sup>-1</sup> (Production of erosion material in the river basin)**  
**R<sub>u</sub> = 0.3034105639269 (Coefficient of the deposit retention)**  
**G<sub>god</sub> = 13157.279085716 m<sup>3</sup> god<sup>-1</sup> (Real soil losses)**  
**G<sub>god</sub> km<sup>-2</sup> = 327.04362723922 m<sup>3</sup> km<sup>-2</sup> god<sup>-1</sup> (Real soil losses per km<sup>2</sup>)**

---

<http://www.wintero.me>