
Web application for Intensity of Erosion and Outflow

Name of the River Basin: Kisjele vode

Country: Montenegro

Year: 2018

**GPS coordinates, latitude and longitude with Google Maps:
43.07919,19.784681**

INPUT DATA

Geometric characteristics of the river basins

F = 10.29929 km² (Surface area of the drainage basin)

O = 14.30003 km (Length of the watershed)

Fv = 7.00498 km² (Surface area of greater portion of the drainage basin)

Fm = 3.29431 km² (Surface area of smaller portion of the drainage basin)

Lv = 4.003 km (Natural length of main water course)

Lb = 5.84308 km (Length of the drainage basin measured by a series of parallel lines)

Topographic characteristics of the river basins

Contour line length - Liz [km]: ["6.10853 ", "7.34426 ", "4.27789 ", "1.76582 ", "1.20790 ", "0.81557 "]

The area between the two neighboring contour lines - f [km²]: ["2.68118 ", "3.11941 ", "2.38172 ", "1.10291 ", "0.50458 ", "0.37513 ", "0.13436 "]

h0 = 600 m (Altitude of the initial contour)

Δh = 100 m (Equidistance)

Hmin = 545 (Lowest altitude in the drainage basin)

Hmax = 1185 (Highest altitude in the drainage basin)

Hydrological characteristics of the river basins

$\Sigma L = 4.003$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

$L_m = 3.89458$ km (The shortest distance between the fountain (head and mouth))

Water permeability

$f_p = 0$ (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0.1833$ (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.8167$ (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

$f_s = 0.390910153$ (Part of the surface area of the drainage basin under the forest)

$f_t = 0.453519217$ (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.155570630$ (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

$h_b = 157.6$ mm (Level of torrent rain)

U_p (years) = 100

$t_o = 8.9$ °C (Average annual air temperature)

$H_{god} = 873.7$ mm (Average annual quantity of precipitation)

Erosion coefficients

$Y = 1.28378$ (Types of soil structures and allied types)

20.42 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

79.58 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

0 % (Epieugleysol and Marshlands)

0 % (Good structured Chernozems and alluvial well-structured deposits)

0 % (Bare, compact igneous)

Xa = 0.53572 (Planning of the drainage basin, rate of drainage basin regulation)

0 % (Bare lands)

15.56 % (Plough-lands)

13.72 % (Orchards and vineyards)

6.91 % (Mountain pastures)

24.72 % (Meadows)

25.41 % (Degraded forests)

13.68 % (Well-constituted forests)

$\phi = 0.28$ (Numerical coefficient of visible and clearly pointed processes of soil erosion)

1.78 % (Depth erosion)

1.58 % (80% of the river basin under rill and gully erosion)

1.38 % (50% of the river basin under rill and gully erosion)

1.19 % (100% of the river basin under surface erosion)

24.72 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)

0.59 % (50% of the river basin under surface erosion)

0.4 % (20% of the river basin under surface erosion)

0 % (There are smaller slides in the watercourse beds)

15.56 % (The river basin mostly under plough-land)

52.8 % (The river basin under forests and perennial vegetation)

INPUT DATA

A = 0.69660400949288 (Coefficient of the river basin form)

m = 0.3518657655853 (Coefficient of the watershed development)

B = 1.7626474393642 km (Average river basin width)

a = 0.72056811683135 ((A)symmetry of the river basin)

G = 0.38866756834694 (Density of the river network of the basin)

K = 1.0278386886391 (Coefficient of the river basin tortuousness)

H_{sr} = 710.05859141747 m (Average river basin altitude)

D = 165.05859141747 m (Average elevation difference of the river basin)

I_{sr} = 20.894615065699 % (Average river basin decline)

H_{leb} = 640 m (The height of the local erosion base of the river basin)

E_r = 113.71775558922 (Coefficient of the erosion energy of the river basins relief)

S₁ = 0.94501 (Coefficient of the regions permeability)

S₂ = 0.7529320954 (Coefficient of the vegetation cover)

W = 1.7229738266463 m (Analytical presentation of the water retention in inflow)

2gDF^{1/2} = 182.63003916662 m km s⁻¹ (Energetic potential of water flow during torrent rains)

Q_{max} = 155.96569211528 m³ s⁻¹ (Maximal outflow from the river basin)

T = 0.99498743710662 (Temperature coefficient of the region)

Z = 0.50694235341003 (Coefficient of the river basin erosion)

W_{god} = 10152.545804758 m³ god⁻¹ (Production of erosion material in the river basin)

R_u = 0.21943022500281 (Coefficient of the deposit retention)

G_{god} = 2227.7754102893 m³ god⁻¹ (Real soil losses)

G_{god} km⁻² = 216.30378504628 m³ km⁻² god⁻¹ (Real soil losses per km²)