
Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S10-1

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.81,57.19

INPUT DATA

Geometric characteristics of the river basins

F = 15.92 km² (Surface area of the drainage basin)

O = 17.41 km (Length of the watershed)

Fv = 8.46 km² (Surface area of greater portion of the drainage basin)

Fm = 7.46 km² (Surface area of smaller portion of the drainage basin)

Lv = 7.3 km (Natural length of main water course)

Lb = 7.18 km (Length of the drainage basin measured by a series of parallel lines)

Topographic characteristics of the river basins

Contour line length - Liz [km]: ["2.33 ", "9.15 ", "13.95 ", "15.71 ", "13.94 ", "10.72 ", "4.53 ", "0.99 "]

The area between the two neighboring contour lines - f [km²]: ["0.34 ", "1.69 ", "2.57 ", "3.50 ", "3.11 ", "3.01 ", "1.43 ", "0.26 ", "0.01 "]

h0 = 900 m (Altitude of the initial contour)

Δh = 100 m (Equidistance)

Hmin = 831 (Lowest altitude in the drainage basin)

Hmax = 1634 (Highest altitude in the drainage basin)

Hydrological characteristics of the river basins

$\Sigma L = 24.83$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

$L_m = 6.46$ km (The shortest distance between the fountain (head and mouth))

Water permeability

$f_p = 0.03$ (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0$ (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.97$ (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

$f_s = 0$ (Part of the surface area of the drainage basin under the forest)

$f_t = 1.00000$ (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.00000$ (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

$h_b = 32.76$ mm (Level of torrent rain)

U_p (years) = 100

$t_o = 12.40$ °C (Average annual air temperature)

$H_{god} = 293.9$ mm (Average annual quantity of precipitation)

Erosion coefficients

$Y = 1.1793$ (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

96.55 % (Decomposed limestone and marls)

0 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

3.45 % (Epieugleysol and Marshlands)

0 % (Good structured Chernozems and alluvial well-structured deposits)

0 % (Bare, compact igneous)

Xa = 0.7 (Planning of the drainage basin, rate of drainage basin regulation)

0 % (Bare lands)

0 % (Plough-lands)

100 % (Orchards and vineyards)

0 % (Mountain pastures)

0 % (Meadows)

0 % (Degraded forests)

0 % (Well-constituted forests)

$\phi = 0.32941$ (Numerical coefficient of visible and clearly pointed processes of soil erosion)

0 % (Depth erosion)

3.56 % (80% of the river basin under rill and gully erosion)

1.61 % (50% of the river basin under rill and gully erosion)

0 % (100% of the river basin under surface erosion)

0 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)

0 % (50% of the river basin under surface erosion)

94.83 % (20% of the river basin under surface erosion)

0 % (There are smaller slides in the watercourse beds)

0 % (The river basin mostly under plough-land)

0 % (The river basin under forests and perennial vegetation)

INPUT DATA

A = 0.46506164383562 (Coefficient of the river basin form)

m = 0.51611489913903 (Coefficient of the watershed development)

B = 2.2172701949861 km (Average river basin width)

a = 0.12562814070352 ((A)symmetry of the river basin)

G = 1.5596733668342 (Density of the river network of the basin)

K = 1.1300309597523 (Coefficient of the river basin tortuousness)

H_{sr} = 1197.6721105528 m (Average river basin altitude)

D = 366.6721105528 m (Average elevation difference of the river basin)

I_{sr} = 44.798994974874 % (Average river basin decline)

H_{leb} = 803 m (The height of the local erosion base of the river basin)

E_r = 127.96167218094 (Coefficient of the erosion energy of the river basins relief)

S₁ = 0.982 (Coefficient of the regions permeability)

S₂ = 0.8 (Coefficient of the vegetation cover)

W = 0.44123541903715 m (Analytical presentation of the water retention in inflow)

2gDF^{1/2} = 338.42307900025 m km s⁻¹ (Energetic potential of water flow during torrent rains)

Q_{max} = 54.555976868649 m³ s⁻¹ (Maximal outflow from the river basin)

T = 1.157583690279 (Temperature coefficient of the region)

Z = 0.82446202617277 (Coefficient of the river basin erosion)

W_{god} = 12737.990108741 m³ god⁻¹ (Production of erosion material in the river basin)

R_u = 0.29209371831941 (Coefficient of the deposit retention)

G_{god} = 3720.686894778 m³ god⁻¹ (Real soil losses)

G_{god} km⁻² = 233.71148836545 m³ km⁻² god⁻¹ (Real soil losses per km²)

<http://www.wintero.me>