
Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S2-intB

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.77,57.63

INPUT DATA

Geometric characteristics of the river basins

F = 10.86 km² (Surface area of the drainage basin)

O = 17.64 km (Length of the watershed)

Fv = 7.75 km² (Surface area of greater portion of the drainage basin)

Fm = 3.1 km² (Surface area of smaller portion of the drainage basin)

Lv = 3.3 km (Natural length of main water course)

Lb = 6.39 km (Length of the drainage basin measured by a series of parallel lines)

Topographic characteristics of the river basins

Contour line length - Liz [km]: ["10.70 ", "16.54 ", "2.57 "]

The area between the two neighboring contour lines - f [km²]: ["3.17 ", "6.56 ", "1.12 ", "0.01 "]

h0 = 1300 m (Altitude of the initial contour)

Δh = 100 m (Equidistance)

Hmin = 1251 (Lowest altitude in the drainage basin)

Hmax = 1565 (Highest altitude in the drainage basin)

Hydrological characteristics of the river basins

$\Sigma L = 18.79$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

$L_m = 3.17$ km (The shortest distance between the fountain (head and mouth))

Water permeability

$f_p = 0.3$ (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0.53$ (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.17$ (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

$f_s = 0$ (Part of the surface area of the drainage basin under the forest)

$f_t = 1.00000$ (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.00000$ (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

$h_b = 33.67$ mm (Level of torrent rain)

U_p (years) = 100

$t_o = 11.70$ °C (Average annual air temperature)

$H_{god} = 304.9$ mm (Average annual quantity of precipitation)

Erosion coefficients

$Y = 0.95085$ (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

70.17 % (Serpentines, red sand stones, flische deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

29.83 % (Epieugleysol and Marshlands)

0 % (Good structured Chernozems and alluvial well-structured deposits)

0 % (Bare, compact igneous)

Xa = 0.6209 (Planning of the drainage basin, rate of drainage basin regulation)

0 % (Bare lands)

0 % (Plough-lands)

20.9 % (Orchards and vineyards)

79.1 % (Mountain pastures)

0 % (Meadows)

0 % (Degraded forests)

0 % (Well-constituted forests)

$\phi = 0.6574$ (Numerical coefficient of visible and clearly pointed processes of soil erosion)

0 % (Depth erosion)

9.13 % (80% of the river basin under rill and gully erosion)

54.53 % (50% of the river basin under rill and gully erosion)

0 % (100% of the river basin under surface erosion)

9.99 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)

0 % (50% of the river basin under surface erosion)

26.35 % (20% of the river basin under surface erosion)

0 % (There are smaller slides in the watercourse beds)

0 % (The river basin mostly under plough-land)

0 % (The river basin under forests and perennial vegetation)

INPUT DATA

A = 1.042363636363636 (Coefficient of the river basin form)

m = 0.28248415675116 (Coefficient of the watershed development)

B = 1.6995305164319 km (Average river basin width)

a = 0.85714285714286 ((A)symmetry of the river basin)

G = 1.7302025782689 (Density of the river network of the basin)

K = 1.0410094637224 (Coefficient of the river basin tortuousness)

H_{sr} = 1338.7348066298 m (Average river basin altitude)

D = 87.7348066298 m (Average elevation difference of the river basin)

I_{sr} = 27.449355432781 % (Average river basin decline)

H_{leb} = 314 m (The height of the local erosion base of the river basin)

E_r = 55.058239303942 (Coefficient of the erosion energy of the river basins relief)

S₁ = 0.661 (Coefficient of the regions permeability)

S₂ = 0.8 (Coefficient of the vegetation cover)

W = 0.46175990646396 m (Analytical presentation of the water retention in inflow)

2gDF^{1/2} = 136.72576933407 m km s⁻¹ (Energetic potential of water flow during torrent rains)

Q_{max} = 34.799843908111 m³ s⁻¹ (Maximal outflow from the river basin)

T = 1.1269427669585 (Temperature coefficient of the region)

Z = 0.69743174994669 (Coefficient of the river basin erosion)

W_{god} = 6827.9692143301 m³ god⁻¹ (Production of erosion material in the river basin)

R_u = 0.18707407508733 (Coefficient of the deposit retention)

G_{god} = 1277.3360254956 m³ god⁻¹ (Real soil losses)

G_{god} km⁻² = 117.61841855392 m³ km⁻² god⁻¹ (Real soil losses per km²)

<http://www.wintero.me>