

---

# Web application for Intensity of Erosion and Outflow

**Name of the River Basin: Shirindareh S7-intC**

**Country: Iran, Islamic Republic of**

**Year: 2019**

**GPS coordinates, latitude and longitude with Google Maps: 37.86,57.36**

## INPUT DATA

### Geometric characteristics of the river basins

**F = 31.89 km<sup>2</sup> (Surface area of the drainage basin)**

**O = 10.99 km (Length of the watershed)**

**Fv = 18.8 km<sup>2</sup> (Surface area of greater portion of the drainage basin)**

**Fm = 13.09 km<sup>2</sup> (Surface area of smaller portion of the drainage basin)**

**Lv = 8.13 km (Natural length of main water course)**

**Lb = 16.57 km (Length of the drainage basin measured by a series of parallel lines)**

### Topographic characteristics of the river basins

**Contour line length - Liz [km]: ["15.24 ", "42.42 ", "22.48 ", "3.93 "]**

**The area between the two neighboring contour lines - f [km<sup>2</sup>]: ["3.59 ", "14.75 ", "11.42 ", "2.01 ", "0.12 "]**

**h0 = 1100 m (Altitude of the initial contour)**

**Δh = 100 m (Equidistance)**

**Hmin = 1029 (Lowest altitude in the drainage basin)**

**Hmax = 1486 (Highest altitude in the drainage basin)**

### Hydrological characteristics of the river basins

---

$\Sigma L = 61.25$  km (The total length of the main watercourse with tributaries of 1<sup>st</sup> and 2<sup>nd</sup> class)

$L_m = 7.19$  km (The shortest distance between the fountain (head and mouth))

### Water permeability

$f_p = 0.23$  (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0$  (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.77$  (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

### Land use

$f_s = 0.03080$  (Part of the surface area of the drainage basin under the forest)

$f_t = 0.95980$  (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.00940$  (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

### Meteorological data

$h_b = 32.68$  mm (Level of torrent rain)

$U_p$  (years) = 100

$t_o = 12.40$  °C (Average annual air temperature)

$H_{god} = 292.9$  mm (Average annual quantity of precipitation)

### Erosion coefficients

$Y = 1.0408$  (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

88.16 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

---

**11.84 % (Epieugleysol and Marshlands)**

**0 % (Good structured Chernozems and alluvial well-structured deposits)**

**0 % (Bare, compact igneous)**

**Xa = 0.60591 (Planning of the drainage basin, rate of drainage basin regulation)**

**0 % (Bare lands)**

**0.94 % (Plough-lands)**

**3.09 % (Orchards and vineyards)**

**92.89 % (Mountain pastures)**

**0 % (Meadows)**

**3.08 % (Degraded forests)**

**0 % (Well-constituted forests)**

**$\phi = 0.5848$  (Numerical coefficient of visible and clearly pointed processes of soil erosion)**

**0 % (Depth erosion)**

**1.32 % (80% of the river basin under rill and gully erosion)**

**54.59 % (50% of the river basin under rill and gully erosion)**

**0 % (100% of the river basin under surface erosion)**

**1.31 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)**

**0 % (50% of the river basin under surface erosion)**

**42.78 % (20% of the river basin under surface erosion)**

**0 % (There are smaller slides in the watercourse beds)**

**0 % (The river basin mostly under plough-land)**

**0 % (The river basin under forests and perennial vegetation)**

## **INPUT DATA**

**A = 0.26359778597786 (Coefficient of the river basin form)**

**m = 0.40612371849772 (Coefficient of the watershed development)**

**B = 1.9245624622812 km (Average river basin width)**

**a = 0.35810598933835 ((A)symmetry of the river basin)**

**G = 1.9206647851991 (Density of the river network of the basin)**

**K = 1.1307371349096 (Coefficient of the river basin tortuousness)**

**H<sub>sr</sub> = 1189.8938538727 m (Average river basin altitude)**

---

---

**D = 160.8938538727 m (Average elevation difference of the river basin)**

**I<sub>sr</sub> = 26.362496080276 % (Average river basin decline)**

**H<sub>leb</sub> = 457 m (The height of the local erosion base of the river basin)**

**E<sub>r</sub> = 61.214273214746 (Coefficient of the erosion energy of the river basins relief)**

**S<sub>1</sub> = 0.862 (Coefficient of the regions permeability)**

**S<sub>2</sub> = 0.79572 (Coefficient of the vegetation cover)**

**W = 0.43875009017647 m (Analytical presentation of the water retention in inflow)**

**2gDF<sup>1/2</sup> = 317.28276993875 m km s<sup>-1</sup> (Energetic potential of water flow during torrent rains)**

**Q<sub>max</sub> = 25.169408247318 m<sup>3</sup> s<sup>-1</sup> (Maximal outflow from the river basin)**

**T = 1.157583690279 (Temperature coefficient of the region)**

**Z = 0.69258698744067 (Coefficient of the river basin erosion)**

**W<sub>god</sub> = 19578.871418555 m<sup>3</sup> god<sup>-1</sup> (Production of erosion material in the river basin)**

**R<sub>u</sub> = 0.14669008660756 (Coefficient of the deposit retention)**

**G<sub>god</sub> = 2872.0263440662 m<sup>3</sup> god<sup>-1</sup> (Real soil losses)**

**G<sub>god</sub> km<sup>-2</sup> = 90.06040589734 m<sup>3</sup> km<sup>-2</sup> god<sup>-1</sup> (Real soil losses per km<sup>2</sup>)**

---

<http://www.wintero.me>